Docket No. ORT-1222 USA DIV

Listing of Claims

Claims 1-27 (Canceled)

28. (Currently Amended) A method of inhibiting platelet aggregation in a subject in need thereof comprising the steps of:

a) selecting a thrombin receptor antagonist compound of Formula II: A method of Claim 27 wherein said compound is selected from a compound of Formula II:

$$X \longrightarrow A_1 \longrightarrow A_2 \longrightarrow Phe$$
 NH₂

wherein:

Y	A ₁	A ₂	<u> </u>
5-(o-Cl-cinnamamido)triazol-3-yl	Cha	Arg	СО
5-(Thien-2-ylacrylamido) triazol-3-yl	Cha	Arg .	CO
5-(Cinnamamido)triazol-3-yl	Cha	Arg	со
5-(α-Me-cinnamamido) triazol-3-yl	Cha	Arg	СО
5-(α-Ph-cinnamamido) triazol-3-yl	Cha	Arg	со
6-Cinnamamidopyridin-3-yl	Cha	Arg	СО
5-Cl, 3-Me-benzothiophen-2-yl	Cha	Arg	SO ₂

Docket No. ORT-1222 USA DIV

Y	<u>A</u> 1	A ₂	<u>x</u>	
5-(p-F-cinnamamido) triazol-3-yl	Cha	Arg	СО	
Benzothiophen-2-yl	Cha	Arg	СО	
1-naphthyl	Cha	Arg	SO ₂	
2-naphthyl	Cha	Атд	SO ₂	-
<u>or</u>				

any pharmaceutically acceptable salt thereof; and

- b) administering said compound to the subject.
- 29. (Currently Amended) A method of Claim 278 wherein said compound is selected from the group consisting essentially of:
- [5-cinnamamidotriazol-3-yl]carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;
- (6-Cinnamamidopyridin-3-yl)carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide; and
- (5-Chloro-3-methyl-benzothiophen-2-yl)carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide.
- 30. (Currently Amended) The method of Claim 278, wherein the therapeutically effective amount of the compound is administered to the subject in an amount from about 0.1 to about 300 mg/kg/day.
- 31. (Currently Amended) The method of Claim 30, wherein the therapeutically effective amount of the compound is about 1 to about 50 mg/kg/day.
- 32. (Cancelled).

Docket No. ORT-1222 USA DIV

33. (Currently Amended) A method of inducing platelet aggregation in a subject in need thereof comprising the steps of:

a) selecting a thrombin receptor antagonist compound of Formula III: A method of Claim 32 wherein said compound is selected from a compound of Formula II:

$$\begin{array}{c}
O \\
\parallel \\
C \\
A_1
\end{array}$$

$$\begin{array}{c}
A_2 \\
A_3
\end{array}$$

$$\begin{array}{c}
NH_2 \\
\end{array}$$

wherein:

Y	<u>A</u> 1	A2	<u>A</u> 3
5-H ₂ N-1,2,4-triazol-3-yl	Cha	Arg	Phe
5-Bromopyridin-3-yl	Cha	Arg	Phe
2-Chromonyl	Cha	Arg	Phe
5-(α-Me-cinnamamido)triazol-3-yl	Cha	Arg	Phe
5-Naphthylacrylamidotriazol-3-yl	Cha	Arg	Phe
Quinoxalin-2-yl	Cha	Arg	Phe
5-(o-Cl-cinnamamido)triazol-3-yl	Cha	Arg	Phe
6-Aminopyridin-3-yl	Cha	Arg	Phe
5-H ₂ N-1,2,4-triazol-3-yl	Cha	Arg	Phe-Arg
Thiadiazol-4-yl	Cha	Arg	Phe

Docket No. ORT-1222 USA DIV

Y	A ₁	A ₂	A ₃
5-(2,3-diMeO-cinnamamido)triazol-3-yl	Cha	Arg	Phe
5-(α-F-cinnamamido)triazol-3-yl	Cha	Arg	Phe
5-(m-NO ₂ -cinnamamido)triazol-3-yl	Cha	Arg	Phe
5-(o-NO₂-cinnamamido)triazol-3-yl	Cha	Arg	Phe
Pyridin-3-yl	Cha	Arg	Phe
5-(m-Cl-cinnamamido)triazol-3-yl	Cha	Arg	Phe
5-H ₂ N-1,2,4-triazol-3-yl	Phe	Arg	Phe
5-H ₂ N-1,2,4-triazol-3-yl	Cha	Lys	Phe
5-H ₂ N-1,2,4-triazol-3-yl	Cha	Arg	Cha
5-H ₂ N-1,2,4-triazol-3-yl	Cha	Arg	Phgly
5-(thiophen-2-ylacrylamido)triazol-3-yl	Cha	Arg	Phe
3-H ₂ N-pyrazin-2-yl	Cha	Arg	Phe
trans 2-(3-pyridyl)ethylenyl	Cha	Arg	Phe
5-(p-MeO-cinnamamido)triazo1-3-yl	Cha	Arg	Phe
5-(p-CN-cinnamamido)triazol-3-yl	Cha	Arg	Phe
5-(p-F-cinnamamido)triazol-3-yl	Cha	Arg	Phe

Docket No. ORT-1222 USA DIV

Y	A ₁	A ₂	A ₃
2-H ₂ N-pyridin-3-yl	Cha	Arg	Phe
5-H ₂ N-1,2,4-triazol-3-yl	Cha	Arg	$ au_{ m yr}$
5-H ₂ N-1,2,4-triazol-3-yl	Cha	Arg	2-Thala
Pyridin-2-yl	Cha	Arg	Phe
5-(p-Phenyl-cinnamamido)triazol-3-yl	Cha	Arg	Phe
N-(p-F-phenylalanyl)-piperidin-3-yl	Cha	Arg	Phe
5-(Cinnamamido)triazol-3-yl	Cha	Агġ	Phe
5-(α-phenyl-cinnamamido)triazo1-3-y1	Cha	Arg	Phe
3-aminophenyl	Cha	Arg	Phe
1-biphenyl	Cha	Arg	Phe
2-biphenylenyl	Cha	Arg	Phe
benzimidazol-5-yl	Cha	Arg	Phe .

and any pharmaceutically acceptable salt thereof; and

- 34. (Currently Amended) A method of Claim 323 wherein said compound is selected from the group consisting essentially of:
- (5-Bromopyridin-3-yl)carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;
- 2-Chromonylcarbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;

b) administering said compound to the subject.

Docket No. ORT-1222 USA DIV

- (5-Aminotriazol-3-yl)carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;
- [5-(\alpha-Methyl)cinnamamidotriazol-3-yl]carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;
- {5-[3-(1-Naphthyl)acrylamido]triazol-3-yl}carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;
- [Quinoxalin-2-yl]carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;
- [5-(o-Chlorocinnamamido)triazol-3-yl]carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;
- (6-Aminopyridin-3-yl)carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;
- (5-Aminotriazol-3-yl)carbonyl-phenylalanyl-arginyl-phenylalanyl-arginineamide;
- (5-Aminotriazol-3-yl)carbonyl-cyclohexylalanyl-lysinyl-phenylalanineamide; and
- {5-[3-(2-Thienyl)acrylamido]triazol-3-yl}carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide.
- 35. (Currently Amended) The method of Claim 323, wherein the therapoutically effective amount of the compound is administered to the subject in an amount from about 0.1 to about 300 mg/kg/day.
- 36. (Currently Amended) The method of Claim 35, wherein the therapeutically effective amount of the compound is about 1 to about 50 mg/kg/day.
- 37. (Cancelled).
- 38. (Currently Amended) A method of Claim 37 wherein said compound is selected from a compound of Formula A method of treating a platelet-mediated thrombotic disorder selected from the group consisting of myocardial infarction, stroke, angina, and ischemic attacks in a subject in need thereof comprising administering to the subject a therapeutically effective amount of a compound of Formula II:

Docket No. ORT-1222 USA DIV

$$X \longrightarrow A_1 \longrightarrow A_2 \longrightarrow NH_2$$
(II)

wherein:

Y	_A ₁	A ₂	_X
5-(o-Cl-cinnamamido)triazol-3-yl	Cha	Arg	со
5-(Thien-2-ylacrylamido) triazol-3-yl	Cha	Arg	со
5-(Cinnamamido)triazol-3-yl	Cha	Arg	СО
5-(α-Me-cinnamamido) triazol-3-yl	Cha	Arg	co
5-(\(\alpha\)-Ph-cinnamamido) triazol-3-yl	Cha	Arg	СО
6-Cinnamamidopyridin-3-yl	Cha	Arg	co
5-Cl, 3-Me-benzothiophen-2-yl	Cha	Arg	SO_2
5-(p-F-cinnamamido) triazol-3-yl	Cha	Arg	CO
Benzothiophen-2-yl	Cha	Arg	CO
1-naphthyl	Cha	Arg	SO ₂
2-naphthyl	Cha	Arg	SO ₂ •


or any pharmaceutically acceptable salt thereof.

Docket No. ORT-1222 USA DIV

- 39. (Currently Amended) A method of Claim 378 wherein said compound is selected from the group consisting essentially of:
- [5-cinnamamidotriazol-3-yl]carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide;
- (6-Cinnamamidopyridin-3-yl)carbonyl-cyclohexylalanyl-arginylphenylalanineamide; and
- (5-Chloro-3-methyl-benzothiophen-2-yl)carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide.
- 40. (Currently Amended) The method of Claim 378, wherein the therapeutically effective amount of the compound is about 0.1 to about 300 mg/kg/day.
- 41. (Previously Presented) The method of Claim 40, wherein the therapeutically effective amount of the compound is about 1 to about 50 mg/kg/day.
- 42. (Cancelled).
- 43. (Currently Amended) A method of Claim 42 wherein said compound is selected from a compound of Formula A method of treating restenosis in a subject in need thereof comprising administering to the subject a therapeutically effective amount of a compound of Formula II:

$$Y \longrightarrow A_1 \longrightarrow A_2 \longrightarrow Phe \longrightarrow NH_2$$

wherein:



Docket No. ORT-1222 USA DIV

Y	Aı	A ₂	х
5-(Thien-2-ylacrylamido) triazol-3-yl	Cha	Arg	СО
5-(Cinnamamido)triazol-3-yl	Cha	Атд	CO
5-(α-Me-cinnamamido) triazol-3-yl	Cha	Arg	CO
5-(α-Ph-cinnamamido) triazol-3-yl	Cha	Arg	СО
6-Cinnamamidopyridin-3-yl	Cha	Arg	со
5-Cl, 3-Me-benzothiophen-2-yl	Cha	Агġ	SO ₂
5-(p-F-cinnamamido) triazol-3-yl	Cha	Arg	со
Benzothiophen-2-yl	Cha	Arg	CO
l-naphthyl	Cha	Arg	SO ₂
2-naphthyl	Cha	Arg	SO ₂

or any pharmaceutically acceptable salt thereof.

- 44. (Currently Amended) A method of Claim 423 wherein said compound is selected from the group consisting essentially of:
- [5-cinnama midotriazol-3-yl] carbonyl-cyclohexylalanyl-arginyl-phenylalanine a mide;
- (6-Cinnamamidopyridin-3-yl)carbonyl-cyclohexylalanyl-arginylphenylalanineamide; and

Docket No. ORT-1222 USA DIV

- (5-Chloro-3-methyl-benzothiophen-2-yl)carbonyl-cyclohexylalanyl-arginyl-phenylalanineamide.
- 45. (Currently Amended) The method of Claim 423, wherein the therapeutically effective amount of the compound is about 0.1 to about 300 mg/kg/day.
- 46. (Previously Presented) The method of Claim 45, wherein the therapeutically effective amount of the compound is about 1 to about 50 mg/kg/day.